arXiv:1202.4204v1 [math.CO] 20 Feb 2012

Vertex Isoperimetric Inequalities for a Family of Graphs on $\mathbb{Z}^k$

Ellen Veomett *
Saint Mary's College of California

A.J. Radcliffe
University of Nebraska-Lincoln

February 21, 2012

Abstract

We consider the family of graphs whose vertex set is $\mathbb{Z}^k$ where two vertices are connected by an edge when their ℓ_∞-distance is 1. We prove the optimal vertex isoperimetric inequality for this family of graphs. That is, given a positive integer n, we find a set $A \subset \mathbb{Z}^k$ of size n such that the number of vertices who share an edge with some vertex in A is minimized. These sets of minimal boundary are nested, and the proof uses the technique of compression.

We also show a method of calculating the vertex boundary for certain subsets in this family of graphs. This calculation and the isoperimetric inequality allow us to indirectly find the sets which minimize the function calculating the boundary.

1 Introduction and Results

For a metric space (X, d) with a notion of volume and boundary, an isoperimetric inequality gives a lower bound on the boundary of a set of fixed volume. Ideally, for any fixed volume, it produces a set of that volume with minimal boundary. The most well-known isoperimetric inequality states that, in Euclidean space, the unique set of fixed volume with minimal boundary is the Euclidean ball.

More recently, questions of isoperimetric inequalities have arisen within various kinds of discrete spaces [12], [7], [9], [11], [10], [16], [20]. Not only are questions regarding isoperimetric inequalities natural geometric questions, but they are known to have many implications in areas such as measure concentration [18], [15], [19], [17] and the theory of random graphs [3], [6], [5], [4].

One of the more broadly known discrete isoperimetric inequalities is Harper's Theorem. Harper's Theorem involves the discrete space $\{0,1\}^n$ and the ℓ_1

---

*The first author was supported for eight weeks during the summer of 2010 through the University of Nebraska-Lincolns Mentoring through Critical Transition Points grant (DMS-0838463) from the National Science Foundation.

metric; that is, the metric defined by the norm

$$||(x_1, x_2, \ldots, x_n)||_1 = \sum_{i=1}^{n} |x_i|$$

A graph $G = (V, E)$ is defined on the vertex set $V = \{0,1\}^n$ where the edge set E consists of points whose distance in the ℓ_1 metric is precisely 1:

$$E = \{(u,v) : u \in \{0,1\}^n, v \in \{0,1\}^n, ||u - v||_1 = 1\}$$

Here, the vertex boundary of a set $A \subset \{0,1\}^n$ consists of all points whose distance from A in the graph metric is no more than 1:

$$\partial A = \{v \in V : d(v, A) \leq 1\}.$$

As usual, the distance between a point x and a set A is defined

$$d(x, A) = \inf_{a \in A} d(x, a) = \inf_{a \in A} \{\text{the length of the shortest path from } x \text{ to } a\}$$

In other words, the vertex boundary consists of both A and all of the neighbors of A. In [12] Harper shows that one can define an ordering on the 0-1 cube $\{0,1\}^n$ such that a set of smallest vertex boundary is achieved on an initial segment. Thus, the sets achieving the minimal boundary are nested.

In [7], Bollobás and Leader found an isoperimetric inequality for a graph which is also defined using the ℓ_1 metric. The vertices of this graph are the vertices of the discrete torus $\mathbb{Z}_m^n$, where $\mathbb{Z}_m$ denotes the integers modulo m and m is necessarily an even integer. The edges of this graph are points whose distance using the ℓ_1 metric is precisely 1. In [7], Bollobás and Leader use a tool called a fractional system to show that the sets of minimal boundary of size $|\{x \in \mathbb{Z}_m^n : ||x - \vec{0}||_1 \leq r\}|$ $(r = 0, 1, 2, \ldots)$ are precisely those balls: $\{x \in \mathbb{Z}_m^n : ||x - \vec{0}||_1 \leq r\}$. Thus, the sets of minimal vertex boundary of size $|\{x \in \mathbb{Z}_m^n : ||x - \vec{0}||_1 \leq r\}|$ $(r = 0, 1, 2, \ldots)$ again are nested.

The fact that sets achieving optimality are nested is crucial in both [12] and [7], as they use the technique of compression. Compression has been utilized in these and many other discrete isoperimetric problems to inductively find sets of minimal boundary, and relies heavily on the fact that the graph and its lower dimensional counterparts have sets of minimal boundary that are nested. Discussions of compression as a technique in discrete isoperimetric problems can be found in [8], [13], [1], and [2].

In the following we consider the vertex isoperimetric inequalities of a family of graphs which was previously unstudied. Specifically, we consider the family of graphs whose vertex set is $\mathbb{Z}^k$ where two vertices are connected by an edge when their ℓ_∞-distance is 1. For each $n \in \mathbb{N}$, we produce a set in $\mathbb{Z}^k$ of size n whose vertex boundary is minimized. We do this by defining a well-ordering $\prec$ on $\mathbb{Z}^k$ and showing that a set of minimal boundary is an initial segment:

Theorem 1. *Let I be an initial segment in $\mathbb{Z}^k$, and A a finite, nonempty subset. If $|I| = |A|$ then $|\partial I| \leq |\partial A|$.*

Thus, there exist sets of minimal boundary which are nested. (The sets of minimal boundary are not unique, as we point out at the end of Section 2). To prove Theorem 1, we use a version of compression which is similar to that used to prove Harper's Theorem in [14]. Compression alone is not enough to produce a set of minimal boundary; instead we show that compressing a set and a particular type of "jostling" eventually results in a set of minimal boundary. We prove Theorem 1 in Section 2.

In Section 3, we show how one can use a "1-dimensional compression" technique to take a set $A \subset \mathbb{Z}^k$ and produce a set of the same size whose boundary is no larger. We call such a set centralized. Then we are able to compute the boundary of centralized sets. This computation along with the isoperimetric inequality of Theorem 1 allows us to deduce which sets achieve the minimum value of the function which computes the boundary of centralized sets. This boundary computation technique may be useful in finding isoperimetric inequalities for other related graphs.

In Section 4 we make some final comments on using the ideas in Section 3 for other graphs.

2 The Proof of Theorem 1

We first consider the graph with vertex set $\mathbb{Z}^k$. Two vertices $x, y \in \mathbb{Z}^k$ are joined by an edge precisely when $\|x-y\|_\infty = 1$. That is, when

$$\max_{i=1,2,\ldots,k} |x_i - y_i| = 1$$

where $x = (x_1, x_2, \ldots, x_k)$ and $y = (y_1, y_2, \ldots, y_k)$.

We define a well-ordering $\prec$ on $\mathbb{Z}^k$ inductively. For $\mathbb{Z}^1$, the well-ordering $\prec$ is:

$$0 \prec 1 \prec -1 \prec 2 \prec -2 \prec 3 \cdots$$

For $k > 1$, the well-ordering $\prec$ on $\mathbb{Z}^k$ is as follows: for $u = (u_1, u_2, \ldots, u_k) \in \mathbb{Z}^k$ define

$$M(u) = \max_{\prec}\{u_i : i = 1, 2, \ldots, k\}$$

where the maximum is according to the previously defined well-ordering on $\mathbb{Z}$. Then for $u, v \in \mathbb{Z}^k$ with $u \neq v$, if $M(u) \prec M(v)$, then $u \prec v$. If $M(u) = M(v)$, define

$$i_u = \min\{i : u_i = M(u) \text{ where } u = (u_1, u_2, \ldots u_k)\}$$
$$i_v = \min\{i : v_i = M(v) \text{ where } v = (v_1, v_2, \ldots v_k)\}$$

If $i_v < i_u$ then $u \prec v$. Finally, if $M(u) = M(v)$ and $i_u = i_v$, we define

$$u' = (u_1, u_2, \ldots, u_{i_u - 1}, u_{i_u + 1}, \ldots u_k) \in \mathbb{Z}^{k-1}$$
$$v' = (v_1, v_2, \ldots, v_{i_u - 1}, v_{i_u + 1}, \ldots v_k) \in \mathbb{Z}^{k-1}$$

and state that $u \prec v$ precisely when $u' \prec v'$.

For example, for $n = 3$, here are the first forty elements according to this well-ordering:

1.$(0,0,0)$
2.$(0,0,1)$
3.$(0,1,0)$
4.$(0,1,1)$
5.$(1,0,0)$
6.$(1,0,1)$
7.$(1,1,0)$
8.$(1,1,1)$
9.$(0,0,-1)$
10.$(0,1,-1)$
11.$(1,0,-1)$
12.$(1,1,-1)$
13.$(0,-1,0)$
14.$(0,-1,1)$
15.$(1,-1,0)$
16.$(1,-1,1)$
17.$(0,-1,-1)$
18.$(1,-1,-1)$
19.$(-1,0,0)$
20.$(-1,0,1)$
21.$(-1,1,0)$
22.$(-1,1,1)$
23.$(-1,0,-1)$
24.$(-1,1,-1)$
25.$(-1,-1,0)$
26.$(-1,-1,1)$
27.$(-1,-1,-1)$
28.$(0,0,2)$
29.$(0,1,2)$
30.$(1,0,2)$
31.$(1,1,2)$
32.$(0,-1,2)$
33.$(1,-1,2)$
34.$(-1,0,2)$
35.$(-1,1,2)$
36.$(-1,-1,2)$
37.$(0,2,0)$
38.$(0,2,1)$
39.$(1,2,0)$
40.$(1,2,1)$

We use the notation $u \preceq v$ to mean that either $u = v$ or $u \prec v$. For $a \in \mathbb{Z}$, we let a^s denote its immediate successor in the well-ordering $\prec$. Thus, for example,

$$\begin{aligned} 1^s &= -1 \\ -1^s &= 2 \\ 0^s &= 1 \end{aligned}$$

Since it will be used later, we present the following remark on how to calculate immediate successors using this well-ordering $\prec$ on $\mathbb{Z}^k$:

Remark 1 (Calculating Successors).

Consider $x = (x_1, x_2, \ldots, x_k) \in \mathbb{Z}^k$. Let m be the smallest entry of x according to the well ordering $\prec$ of $\mathbb{Z}$. Thus,

$$m = x_i \text{ for some } i$$

and

$$m \preceq x_i \text{ for each } i$$

Define

$$i_m = \max\{i : x_i = m\}.$$

If $i_m = k$, then $x_1 = x_2 = \ldots = x_k$ and the immediate successor of x is $(0, 0, \ldots, 0, x_k^s)$.

Otherwise, $i_m < k$. In this case, the immediate successor to x is the vector $x^s = (y_1, y_2, \ldots, y_k)$ where x^s has the same entries as x except that all entries x_j which are equal to x_{i_m} with $j < i_m$ change to 0 (or stay 0 if $x_{i_m} = 0$), x_{i_m} changes to $x_{i_m}^s$, and all entries x_j which are equal to $x_{i_m}^s$ with $j > i_m$ change to 0. That is,

$$y_i = \begin{cases} 0 & \text{if } i < i_m \text{ and } x_i = x_{i_m} \text{ or if } i > i_m \text{ and } x_i = x_{i_m}^s \\ x_{i_m}^s & \text{if } i = i_m \\ x_i & \text{otherwise} \end{cases}$$

For a set $A \subset \mathbb{Z}^k$, we use the notation ∂A to denote

$$\partial A = \{x \in \mathbb{Z}^k : \|x - a\|_\infty \leq 1 \text{ for some } a \in A\}.$$

Thus, $A \subset \partial A$. We note that this is the notation used in [14], [9], [10], and others. We use $|A|$ to denote the size of a finite set.

Our goal will be to prove that, for any $A \subset \mathbb{Z}^k$ $|\partial A| \geq |\partial I_{|A|}|$ where $I_{|A|}$ is the initial segment in $\mathbb{Z}^k$ of length $|A|$, according to the well-ordering $\prec$. In order to prove this, we will need some Lemmas about initial segments in $\mathbb{Z}^k$ and their boundaries.

For a number $a \in \mathbb{Z}$, we let a^+ denote the element of $\{a, a+1, a-1\}$ which is largest in the well-ordering $\prec$ and a^- the element of $\{a, a+1, a-1\}$ which is the smallest. Thus, for example,

$$\begin{aligned} 1^+ &= 2 & \qquad 1^- &= 0 \\ -1^+ &= -2 & \qquad -1^- &= 0 \\ 0^+ &= -1 & \qquad 0^- &= 0 \end{aligned}$$

Lemma 1. *If $I \subset \mathbb{Z}^k$ is an initial segment, so is ∂I*

Proof. By induction on $|I|$. If $|I| = 1$, then $I = \{(0, 0, \ldots, 0)\}$ and $\partial I = \{0, 1, -1\}^k$ which is the initial segment of size 3^k. Now let $v \in \mathbb{Z}^k$, and suppose that $I = \{x \in \mathbb{Z}^k : x \preceq v\}$ and ∂I are initial segments. Let u be the successor of v. Note that u is not the zero vector. We will show that $\partial(I \cup \{u\})$ is an initial segment.

Let $v = (v_1, v_2, \ldots, v_k)$. Note that the vector in $\mathbb{Z}^k$ which is an element of ∂I and the latest in the well-ordering $\prec$ is $(v_1^+, v_2^+, \ldots, v_k^+)$. Thus, by our inductive assumption,

$$\partial I = \{x \in \mathbb{Z}^k : x \preceq (v_1^+, v_2^+, \ldots, v_k^+)\}.$$

Note that

$$\partial(I \cup \{u\}) = \partial I \cup \{(u_1 + \delta_1, u_2 + \delta_2, \ldots, u_k + \delta_k) : (\delta_1, \delta_2, \ldots, \delta_k) \in \{0, 1, -1\}^k\}.$$

Let $i \in \{1, 2, \ldots, k\}$ and suppose that $u_i \neq 0$. Then

$$(u_1, u_2, \ldots, u_{i-1}, u_i^-, u_{i+1}, \ldots, u_n) \prec (u_1, u_2, \ldots, u_n)$$

and for $(\delta_1, \delta_2, \ldots, \delta_{k-1}) \in \{0, 1, -1\}^{k-1}$ each of

$$\begin{aligned} &(u_1 + \delta_1, u_2 + \delta_2, \ldots, u_{i-1} + \delta_{i-1}, u_i^-, u_{i+1} + \delta_i, \ldots, u_k + \delta_{k-1}) \\ &(u_1 + \delta_1, u_2 + \delta_2, \ldots, u_{i-1} + \delta_{i-1}, u_i, u_{i+1} + \delta_i, \ldots, u_k + \delta_{k-1}) \end{aligned}$$

are adjacent to $(u_1, u_2, \ldots, u_{i-1}, u_i^-, u_{i+1}, \ldots, u_n)$.

Thus, the elements $(x_1, x_2, \ldots, x_n)$ of $\partial(I \cup \{u\})$ which are *not* in ∂I are precisely those of the form:

$$x_i = \begin{cases} u_i^+ & \text{if } u_i \neq 0 \\ 0, 1, \text{ or } -1 & \text{if } u_i = 0 \end{cases}$$

Note that if $u_i \neq 0$ then u_i^+ is at least as large as 2 in the well-ordering $\prec$ on $\mathbb{Z}$. Finally, recall that u is the immediate successor to $v = (v_1, v_2, \ldots, v_k)$ and that every entry of $(v_1^+, v_2^+, \ldots, v_k^+)$ is at least as large as -1 in the well-ordering on $\mathbb{Z}$. Thus, by Remark 1, we can see that the elements of $\partial(I \cup \{u\})$ which are *not* in ∂I are the 3^ℓ immediate successors of $(v_1^+, v_2^+, \ldots, v_k^+)$ where ℓ is the number of 0 entries of u.

$\square$

We note that we have also proved the following:

Lemma 2. *Suppose I is an initial segment in $\mathbb{Z}^k$ and let v be the first element not in I. Then*

$$|\partial(I \cup \{v\})| = |\partial I| + 3^\ell$$

where ℓ is the number of coordinates equal to 0 in v.

The following technical Lemma will be used in the proof of Theorem 1

Lemma 3. *Suppose $k \geq 2$ and consider a segment in $\mathbb{Z}^k$:*

$$S = \{v \in \mathbb{Z}^k : v_0 \leq v \leq v_1\}, \quad k \geq 2.$$

Let $v_0 = (v_{0,1}, v_{0,2}, \ldots, v_{0,k})$ and suppose that $v_{0,j} = 0$ for all $j \neq j^*$ and $v_{0,j^*} \neq 0$. Suppose that the last t vectors in S all do not have a 0 entry. Then for any $j \in \{1, 2, \ldots, k\}$, $j \neq j^*$ and for any

$$s \in \left\{1, -1, 2, -2, \ldots (-1)^{t+1} \left\lceil \frac{t}{2} \right\rceil \right\}$$

there is a vector in S whose jth entry is s.

Proof. Note that the numbers in the set $\left\{1, -1, 2, -2, \ldots (-1)^{t+1} \left\lceil \frac{t}{2} \right\rceil \right\}$ are the first t elements in the list $1, -1, 2, -2, 3, -3, \ldots$.

Recall from Remark 1 how immediate successors in $\mathbb{Z}^k$ are calculated. The only way that t successive vectors all do not have a 0 entry is if a single coordinate is increasing (according to $\prec$) in those successive vectors.

Suppose it is the ith coordinate which is increasing in the final t vectors of S. This implies that, in *each* of the t successive vectors that have no 0 coordinate, the ith coordinate is smaller than any of the other coordinates (in the $\prec$ ordering on $\mathbb{Z}$). Thus, for *any* $\ell \neq i$, the ℓth coordinate must be at least as large as the tth element in the list $1, -1, 2, -2, \ldots$. Hence we can see that every coordinate of v_1 has a value at least as large as the tth element in the list $1, -1, 2, -2, 3, \ldots$.

Consider any $j \neq j^*$. By assumption, the jth coordinate v_0 is 0. Recall that every coordinate of v_1 has a value at least as large as the tth element in the list $1, -1, 2, -2, 3, \ldots$ and S is the segment between v_0 and v_1. Thus, by the definition of the ordering $\prec$ on $\mathbb{Z}^k$, for the 1st through tth elements in the list $1, -1, 2, -2, \ldots$, there must be an element between v_0 and v_1 whose jth coordinate is equal to that element.

Specifically, let $v_0 = (v_{0,1}, v_{0,2}, \dots, v_{0,k})$ and $s \in \{1, -1, 2, \dots, (-1)^{t+1}\left\lceil \frac{t}{2} \right\rceil\}$. Then the vector $x = (x_1, x_2, \dots, x_k)$ where

$$x_\ell = \begin{cases} v_{0,\ell} & \text{if } \ell \neq j \\ s & \text{if } \ell = j \end{cases}$$

has an entry equal to s and $v_0 \prec x \preceq v_1$. Thus, we have proven the Lemma.

□

The notation we use (and the technique used) in proving Theorem 1 is similar to that in that in the proof of Harper's Theorem in [14]. For $A \subset \mathbb{Z}^k$, $i \in \{1, 2, \dots, k\}$, and $j \in \mathbb{Z}$ we define A_{i^j} to be the set of vectors in $\mathbb{Z}^{k-1}$ such that, when a j is inserted in between the $i-1$th and ith entries, the resulting vector is in A. That is,

$$A_{i^j} = \{x \in \mathbb{Z}^{k-1} : (x_1, x_2, \dots, x_{i-1}, j, x_i, x_{i+1}, \dots, x_{k-1}) \in A\}$$

Now we are ready to prove Theorem 1.

Proof of Theorem 1. Let $A \subset \mathbb{Z}^k$ be finite and nonempty. We proceed by induction on k. For $k = 1$, one can easily see that

$$|\partial A| = |A| + 2$$

if A is a segment: $A = \{j \in \mathbb{Z} : a \leq j \leq b \text{ for some } a, b \in \mathbb{Z}\}$ and

$$|\partial A| > |A| + 2$$

if A is not a segment. Since every initial segment in $\mathbb{Z}$ according to the well-ordering $\prec$ is a segment, the Theorem is proved for $k = 1$.

Now suppose $A \subset \mathbb{Z}^k$ is finite and nonempty, $k > 1$, and the theorem is proven for all smaller positive k. Let $i \in \{1, 2, \dots, k\}$. Note that there are only finitely many $j \in \mathbb{Z}$ for which the set

$$A_{i^j}$$

is nonempty, and that

$$\sum_{j \in \mathbb{Z}} |A_{i^j}| = |A|$$

We define a set C_i which is the "i-compression" of A by specifying its $k-1$-dimensional sections fixing the ith coordinate. Specifically, for $j \in \mathbb{Z}$, we define $(C_i)_{i^j}$ to be the initial segment in $\mathbb{Z}^{k-1}$ of size $|A_{i^j}|$. This gives $|(C_i)_{i^j}| = |A_{i^j}|$ for each j. Thus, since

$$\sum_{j \in \mathbb{Z}} |A_{i^j}| = |A|$$

and

$$\sum_{j \in \mathbb{Z}} |(C_i)_{i^j}| = |C_i|$$

we have $|C_i| = |A|$. Now note that

$$\begin{aligned}(\partial A)_{i^j} &= \partial(A_{i^j}) \cup \partial(A_{i^{j-1}}) \cup \partial(A_{i^{j+1}}) \\ (\partial C_i)_{i^j} &= \partial((C_i)_{i^j}) \cup \partial((C_i)_{i^{j-1}}) \cup \partial((C_i)_{i^{j+1}})\end{aligned}$$

By definition, we know that $|(C_i)_{i^j}| = |A_{i^j}|$, $|(C_i)_{i^{j-1}}| = |A_{i^{j-1}}|$, and $|(C_i)_{i^{j+1}}| = |A_{i^{j+1}}|$. Also, $(C_i)_{i^j}$ is an initial segment, so by induction we have $|\partial((C_i)_{i^j})| \leq |\partial(A_{i^j})|$. Similarly for $j+1$ and $j-1$.

If the union above were necessarily disjoint, we would be done. Since it's not, we note the following: all of $\partial((C_i)_{i^j})$, $\partial((C_i)_{i^{j-1}})$ and $\partial((C_i)_{i^{j+1}})$ are initial segments. Thus, they are ordered by containment. Thus, either the inequality $|\partial((C_i)_{i^j})| \leq |\partial(A_{i^j})|$ or one of the equalities $|\partial((C_i)_{i^{j-1}})| \leq |\partial(A_{i^{j-1}})|$, $|\partial((C_i)_{i^{j+1}})| \leq |\partial(A_{i^{j+1}})|$ is enough to give us that $|(\partial C_i)_{i^j}| \leq |(\partial A)_{i^j}|$.

Note that $j \in \mathbb{Z}$ was arbitrary. Thus, since

$$\begin{aligned}\sum_{j\in\mathbb{Z}} |(\partial A)_{i^j}| &= |\partial A| \\ \sum_{j\in\mathbb{Z}} |(\partial C_i)_{i^j}| &= |\partial C_i|\end{aligned}$$

we have $|\partial C_i| \leq |\partial A|$.

Hence, we have shown that, for $i \in \{1, 2, \ldots, k\}$, the ith-compression of A is a set of the same size with no larger boundary. We note that a set which is compressed in every coordinate need not be an initial segment. For example, for $k = 3$, the set

$$\{(0,0,0), (0,0,1), (0,1,0), (1,0,0)\}$$

is i-compressed, but not an initial segment.

Let $X \subset \mathbb{Z}^k$ be a set of size $|A|$ which has the following properties:

1. $|\partial X| \leq |\partial A|$
2. X is compressed in every coordinate $i \in \{1, 2, \ldots, k\}$.
3. If Z is any other set of size $|A|$ satisfying the above properties 1 and 2, let $\ell(X)$ and $\ell(Z)$ be the last elements of X and Z in the well-ordering $\prec$ respectively. Then we must have

$$\ell(X) \preceq \ell(Z)$$

In words: X is a set of size $|A|$ with boundary no larger than the boundary of A which is compressed in every coordinate (the previous arguments show that such an X exists). We also choose X such that it is "as close as possible to an initial segment" as described in property 3: its last element is as small as possible. We will show that X must be an initial segment by contradiction.

Suppose that X is not an initial segment. Then there exists $x \notin X$, $y \in X$, such that $x \prec y$. We note that, if $x_i = y_i$ for some $i \in \{1, 2, \ldots, k\}$, then defining

$$x^* = (x_1, \ldots, x_{i-1}, x_{i+1}, \ldots, x_k)$$
$$y^* = (y_1, \ldots, y_{i-1}, y_{i+1}, \ldots, y_k)$$

we have $x^* \prec y^*$ in $\mathbb{Z}^{k-1}$, contradicting the fact that X is i-compressed. Thus, each of the coordinates of x and y must be different.

Let S_α be the first segment in X. That is, $S_\alpha \subset X$, S_α is a segment according to the ordering $\prec$, and the immediate successor to the last element in S_α is not in X. Note that, since X is compressed in every coordinate, this implies that $(0, 0, \ldots, 0) \in S_\alpha$. Let y be the first element in $X \backslash S_\alpha$, and let x be its immediate predecessor. From the above comments, we know that either

$$x = (0, 1, 1, \ldots, 1) \quad y = (1, 0, 0, \ldots, 0)$$

or $x_1 \neq 0$ and

$$x = (\underbrace{x_1, x_1, \ldots, x_1}_{\ell}, \underbrace{x_1^s, x_1^s, \ldots, x_1^s}_{n-\ell}) \quad y = (\underbrace{0, 0, \ldots, 0}_{\ell-1}, x_1^s, \underbrace{0, 0, \ldots, 0}_{k-\ell})$$

Case 1:

$$x = (0, 1, 1, \ldots, 1) \quad y = (1, 0, 0, \ldots, 0)$$

Let S_Ω be the last segment in X. That is, $S_\Omega = \{w \in \mathbb{Z}^k : u \preceq w \preceq v\}$, $S_\Omega \subset X$, v is the last element in X, and the immediate predecessor to u is not in X. Since $x \notin X$ and $u \in X$, by the above arguments we must have $u_1 \neq 0$ and

$$u = (u_1, \underbrace{0, 0, \ldots, 0}_{k-1})$$

for some $u_1 \neq 0$. Additionally, we note that the successor to u is $(u_1, 0, \ldots, 0, 1)$ which must have a coordinate in common with x. Thus, by the above arguments, we must have $u = v$. We note that by Lemma 2, there are at least 3^{k-1} elements which adjacent to u and are not adjacent to any other element in X. There are 3 elements which are adjacent to x and not adjacent to any element of $S_\alpha \subset X$. Thus, we can see that

$$\partial((X \cup \{x\}) \backslash \{u\}) \leq \partial(X) - 3^{k-1} + 3 \leq \partial(X), \quad |(X \cup \{x\}) \backslash \{u\}| = |X|$$

and the set $(X \cup \{x\}) \backslash \{u\}$ is closer to an initial segment with respect to property 3 above. Thus, we have reached a contradiction in Case 1.

Case 2: $x_1 \neq 0$ and

$$x = (\underbrace{x_1, x_1, \ldots, x_1}_{\ell}, \underbrace{x_1^s, x_1^s, \ldots, x_1^s}_{k-\ell}) \quad y = (\underbrace{0, 0, \ldots, 0}_{\ell-1}, x_1^s, \underbrace{0, 0, \ldots, 0}_{k-\ell})$$

Let S_Ω be the last segment in X. That is, $S_\Omega = \{w \in \mathbb{Z}^k : u \preceq w \preceq v\}$, $S_\Omega \subset X$, v is the last element in X, and the immediate predecessor to u is not in X.

Let S_β be the first segment not in X. That is, $S_\beta = \{z \in \mathbb{Z}^k : q \preceq z \preceq x\}$ and the immediate predecessor of q is the last element of S_α. Note that, by the above arguments, since $q \notin X$ and $y \in X$ all of the entries of q and y must be different. Thus, the vector q is somewhere in between

$$\begin{cases} (\underbrace{x_1, \dots, x_1}_{\ell-2}, 1, x_1, \underbrace{x_1^s, \dots, x_1^s}_{k-\ell}) \text{ and } x = (\underbrace{x_1, \dots, x_1}_{\ell}, \underbrace{x_1^s, \dots, x_1^s}_{k-\ell}) & \text{if } 1 < \ell < k \\ (x_1, x_1^s, \dots, x_1^s, 1) \text{ and } x = (x_1, x_1^s, x_1^s, \dots, x_1^s) & \text{if } \ell = 1 \\ (x_1, \dots, x_1, 1) \text{ and } x = (x_1, x_1, \dots, x_1) & \text{if } \ell = k \end{cases}$$

If q has no 0 entries, then we know that there is no more than $3^0 = 1$ element in $\mathbb{Z}^k$ adjacent to q which is not adjacent to any other element of $S_\alpha \subset X$. Also there are $3^t \geq 1$ elements in $\mathbb{Z}^k$ larger than v which are adjacent to v and not adjacent to any other element in X, where t is equal to the number of 0 entries of v. Thus, we have

$$\partial((X \cup \{q\}) \backslash \{v\}) \leq \partial(X) - 3^t + 1 \leq \partial(X), \quad |(X \cup \{q\}) \backslash \{v\}| = |X|$$

and $(X \cup \{q\}) \backslash \{v\}$ is closer to an initial segment with respect to property 3 above.

It is possible for q to have a 0 entry. Specifically, we could have

$$q = (\underbrace{x_1, \dots, x_1}_{\ell-1}, 0, \underbrace{x_1^s, \dots, x_1^s}_{k-\ell}) \quad \text{if } 1 < \ell < k$$

or we could have

$$q = (0, x_1^s, \dots, x_1^s) \quad \text{if } \ell = 1$$

If q does have a 0 entry, then we exchange for q instead of v, we exchange for the immediate successor of q instead of the immediate predecessor of v, and so on until either we have exchanged out all of S_Ω or filled all of S_β. This creates a new X' of the same size as X. Note that q is the only element of S_β which has a 0 entry. We claim that at least one of the elements of S_Ω that we exchanged for an element of S_β had a 0 entry. Indeed, if all of S_Ω is exchanged, we know that u has a 0 entry. Otherwise, only the last t entries of S_Ω were exchanged, where x_1 is the tth entry in the list $0, 1, -1, 2, -2, \dots$. If none of them had a 0 entry, then Lemma 3 would imply that one of the vectors in S_Ω has an entry equal to x_1^s and another vector in S_Ω has that same entry equal to x_1. This contradicts the fact that no element of S_Ω can share any entries with x.

Thus, at least one of the elements of S_Ω that we exchanged for an element of S_β had a 0 entry. Thus, by Lemma 2, we know that the boundary of the new set X' is no larger than the boundary of X and that X' is closer to an initial segment with respect to property 3 above. Thus, we have reached a contradiction in Case 2.

Hence, X must indeed be an initial segment, and we have proved our Theorem.

□

We note that the sets that we found here of minimum boundary are not unique. For example, the sets in Figure 2 are both sets in $\mathbb{Z}^2$ of size 10 of minimal boundary, but they are not isomorphic. The blue dots represent the set, while the red dots represent the additional vertices which are in the boundary of the set.



Figure 1: The set on the left is an initial segment, while the set on the right is not.

Finally, we note that the same arguments utilized above can also be used to prove a similar isoperimetric inequality for $\mathbb{N}^k$. That is, we consider the graph whose vertex set is $\mathbb{N}^k$ such that there is an edge between $x \in \mathbb{N}^k$ and $y \in \mathbb{N}^k$ precisely when $\|x - y\|_\infty = 1$. A well-ordering $\prec_{\mathbb{N}}$ is defined on $\mathbb{N}^k$ inductively just as it is for $\mathbb{Z}^k$, but with the base case of $\mathbb{N}^1$ being the standard ordering of $\mathbb{N}$:

$$0 < 1 < 2 < 3 < \cdots$$

All of the Lemmas and the Theorem are proven similarly. In the case of $\mathbb{N}^k$, the equation of Lemma 2 would instead conclude that

$$|\partial(I \cup \{v\})| = |\partial I| + 2^\ell$$

where ℓ is the number of coordinates equal to 0 in v. With this well-ordering $\prec_{\mathbb{N}}$, we have the following:

Corollary 1. *Let I be an initial segment in $\mathbb{N}^k$, and A a finite, nonempty subset. If $|I| = |A|$ then $|\partial I| \leq |\partial A|$.*

3 Boundary Computations

We define a graph on the vertex set $\mathbb{Z}^k \times \mathbb{N}^d$, where k and d are non-negative integers (not both 0). We use $\mathbb{N}$ to denote the set of non-negative integers:

$$\mathbb{N} = \{x \in \mathbb{Z} : x \geq 0\}$$

As usual, we say that two vertices $x, y \in \mathbb{Z}^k \times \mathbb{N}^d$ are joined by an edge precisely when their ℓ_∞-distance is 1. That is, when

$$\|x - y\|_\infty = \max_{i=1,2,\ldots,k+d} \{|x_i - y_i|\} = 1.$$

For simplicity, we introduce the following notation: for a real-valued vector $p = (p_1, p_2, \ldots, p_n) \in \mathbb{R}^n$, and $x \in \mathbb{R}$, we define

$$(p, x \to i) = (p_1, p_2, \ldots, p_{i-1}, x, p_i, p_{i+1}, \ldots, p_n) \in \mathbb{R}^{n+1}$$

In words, $(p, x \to i)$ is the vector that results when placing x in the ith coordinate of p and shifting the ith through nth coordinates of p to the right.

There are two types of 1-dimensional compression which we will utilize.

Definition 1. *We say that a set $S \subset \mathbb{Z}^k \times \mathbb{N}^d$ is* centrally compressed *in the i-th coordinate ($1 \leq i \leq k$) with respect to $p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d$ if the set*

$$\{x \in \mathbb{Z} : (p, x \to i) \in S\}$$

is either empty or of one of the following two forms:

$$\{x : -a \leq x \leq a \text{ for } a \in \mathbb{N}\}$$
$$OR$$
$$\{x : -a \leq x \leq a + 1 \text{ for } a \in \mathbb{N}\}$$

Definition 2. *We say that $S \subset \mathbb{Z}^k \times \mathbb{N}^d$ is* downward compressed *in the j-th coordinate ($k + 1 \leq j \leq k + d$) with respect to $p \in \mathbb{Z}^k \times \mathbb{N}^{d-1}$ if the set*

$$\{x \in \mathbb{N} : (p, x \to j) \in S\}$$

is either empty or of the form:

$$\{x : 0 \leq x \leq a \text{ for } a \in \mathbb{N}\}$$

For certain subsets $S \subset \mathbb{Z}^k \times \mathbb{N}^d$, we will calculate the boundary

$$\begin{aligned} \partial S &= \{x \in \mathbb{Z}^k \times \mathbb{N}^d : \|y - x\|_\infty \leq 1 \text{ for some } y \in S\} \\ &= \left\{x \in \mathbb{Z}^k \times \mathbb{N}^d : \exists\, s \in S \text{ such that } x = s + \epsilon \text{ for some } \epsilon \in \{-1, 0, 1\}^{k+d}\right\} \end{aligned}$$

Specifically, we will calculate the boundary of sets which are centrally compressed in the ith coordinate with respect to any $p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d$ for $1 \leq i \leq k$, and which are downward compressed in the jth coordinates with respect to any $p \in \mathbb{Z}^k \times \mathbb{N}^{d-1}$ for $k + 1 \leq j \leq k + d$. We will say that such sets are "compressed in every coordinate." Before completing the calculations, we shall see that a set of minimum boundary must be compressed in every coordinate.

3.1 Effect of Compression

Definition 3. *Let $S \subset \mathbb{Z}^k \times \mathbb{N}^d$. For $1 \leq i \leq k$, we define S_i to be the ith central compression of S by specifying its 1-dimensional sections in the ith coordinate. Specifically,*

1. *For each $p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d$,*

$$|\{x \in \mathbb{Z} : (p, x \to i) \in S_i\}| = |\{x \in \mathbb{Z} : (p, x \to i) \in S\}|$$

2. *S_i is centrally compressed in the ith coordinate with respect to p for each $p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d$.*

For $k+1 \leq j \leq k+d$, we define S_j to be the jth central compression of S by specifying its 1-dimensional sections in the jth coordinate. Specifically,

1. *For each $p \in \mathbb{Z}^k \times \mathbb{N}^{d-1}$,*

$$|\{x \in \mathbb{N} : (p, x \to j) \in S_j\}| = |\{x \in \mathbb{N} : (p, x \to j) \in S\}|$$

2. *S_j is downward compressed in the jth coordinate with respect to p for each $p \in \mathbb{Z}^k \times \mathbb{N}^{d-1}$.*

In words: after fixing a coordinate $i \in \{1, 2, \ldots, k\}$, we consider all lines in $\mathbb{Z}^k \times \mathbb{N}^d$ where only the ith coordinate varies, and we intersect those lines with S. Eeach of the points in those intersections are moved along the line so that they are a segment centered around 0. The result is S_i. Similarly, after choosing $j \in \{k+1, \ldots, k+d\}$, we again consider lines in $\mathbb{Z}^k \times \mathbb{N}^d$ where the jth coordinate varies, and intersect those lines with S. We now move the points along the line so that they are a segment starting from 0. The result is S_j.

Proposition 1. *Suppose that $S \subset \mathbb{Z}^k \times \mathbb{N}^d$. For $1 \leq i \leq k$, let S_i be the ith central compression of S. Then*

$$\partial S_i \leq \partial S$$

Similarly, for $k+1 \leq j \leq k+d$, let S_j be the jth downward compression of S. Then

$$\partial S_j \leq \partial S$$

Proof. Let $1 \leq i \leq k$ and fix $p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d$. Consider any $\epsilon \in \{-1, 0, 1\}^{k+d-1}$ for which

$$p + \epsilon \in \mathbb{Z}^{k-1} \times \mathbb{N}^d.$$

Let

$$I = \{\epsilon \in \{-1, 0, 1\}^{k+d-1} : \exists\, x \in \mathbb{Z} : (p + \epsilon, x \to i) \in S\}.$$

Note that, by the definition of S_i, for $p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d$, we have

$$\{x \in \mathbb{Z} : (p + \epsilon, x \to i) \in S\} \neq \emptyset$$

if and only if

$$\{x \in \mathbb{Z} : (p + \epsilon, x \to i) \in S_i\} \neq \emptyset$$

We have

$$\begin{aligned}
&\{(p, x \to i) : x \in \mathbb{Z}\} \cap \partial S \\
&\qquad = \bigcup_{\epsilon \in I} \{(p, x \to i) : \exists\, y \in \{x-1, x, x+1\} \text{such that } (p + \epsilon, y \to i) \in S\}
\end{aligned}$$

Thus, we have

$$\begin{aligned}
&|\{(p, x \to i) : x \in \mathbb{Z}\} \cap \partial S| \\
&\geq \max_{\epsilon \in I} |\,\{(p, x \to i) : \exists\, y \in \{x-1, x, x+1\}\text{such that } (p+\epsilon, y \to i) \in S\}\,| \\
&\geq \max_{\epsilon \in I} |\{x \in \mathbb{Z} : (p+\epsilon, x \to i) \in S\}| + 2
\end{aligned}$$

where the last inequality is achieved if and only if, for the ϵ achieving the maximum,

$$\{x \in \mathbb{Z} : (p+\epsilon, x \to i) \in S\} = \{x \in \mathbb{Z} : a \leq x \leq b\}$$

for some $a, b \in \mathbb{Z}$.

Additionally, we note that

$$\begin{aligned}
&\{(p, x \to i) : x \in \mathbb{Z}\} \cap \partial S_i \\
&\quad = \bigcup_{\epsilon \in I} \{(p, x \to i) : \exists\, y \in \{x-1, x, x+1\} \text{ such that } (p+\epsilon, y \to i) \in S_i\}\,.
\end{aligned}$$

so that now, by the definition of S_i,

$$\begin{aligned}
&|\{(p, x \to i) : x \in \mathbb{Z}\} \cap \partial S_i| \\
&= \max_{\epsilon \in I} |\,\{(p, x \to i) : \exists\, y \in \{x-1, x, x+1\} \text{ such that } (p+\epsilon, y \to i) \in S_i\}\,| \\
&= \max_{\epsilon \in I} |\{x \in \mathbb{Z} : (p+\epsilon, x \to i) \in S_i\}| + 2
\end{aligned}$$

By the definition of S_i, for each $\epsilon \in I$, we have

$$|\{x \in \mathbb{Z} : (p+\epsilon, x \to i) \in S\}| = |\{x \in \mathbb{Z} : (p+\epsilon, x \to i) \in S_i\}|.$$

Finally, we note that both ∂S and ∂S_i are the disjoint unions of those 1-dimensional sections:

$$\begin{aligned}
\partial S &= \bigcup_{p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d} \{(p, x \to i) : x \in \mathbb{Z}\} \cap \partial S \\
\partial S_i &= \bigcup_{p \in \mathbb{Z}^{k-1} \times \mathbb{N}^d} \{(p, x \to i) : x \in \mathbb{Z}\} \cap \partial S_i
\end{aligned}$$

Thus, we can conclude that

$$\partial S \geq \partial S_i.$$

The proof in the case of downward compression in coordinate $j \in \{k+1, \ldots, k+d\}$ is similar. $\square$

3.2 Boundaries of Compressed Subsets of $\mathbb{Z}^k \times \mathbb{N}^d$

We have the following definitions: For fixed $k, d \in \mathbb{N}$ (not both 0), let

$$\begin{aligned} K &= \{1, 2, \ldots, k\} \\ D &= \{k+1, k+2, \ldots, k+d\} \end{aligned}$$

In addition, for $I \subset \{1, 2, \ldots, k+d\}$ and $S \subset \mathbb{Z}^k \times \mathbb{N}^d$, let $P_I(S)$ denote the $k + d - |I|$-dimensional projection of S which results from deletion of the coordinates in I.

Theorem 2. *Let $S \subset \mathbb{Z}^k \times \mathbb{N}^d$ be a finite set which is centrally compressed in coordinate i for each $1 \leq i \leq k$ and downward compressed in each coordinate j where $k+1 \leq j \leq k+d$. Then*

$$\partial S = \sum_{I \subset \{1,2,\ldots,k+d\}} 2^{|I \cap K|} |P_I(S)|.$$

Proof. We proceed by induction on $|S|$. If $|S| = 1$, then $S = \{(0, 0, 0, \ldots, 0)\}$ and

$$\begin{aligned} \partial S = \{(x_1, x_2, \ldots, x_{k+d}) : x_i \in \{-1, 0, 1\} \text{ for } 1 \leq i \leq k, \\ x_j \in \{0, 1\} \text{ for } k+1 \leq j \leq k+d\} \end{aligned}$$

so that

$$|\partial S| = 3^k 2^d$$

We note that in this case, $|P_I(S)| = 1$ for any $I \subset \{1, 2, \ldots, k+d\}$ so that

$$\begin{aligned} \sum_{I \subset \{1,2,\ldots,k+d\}} 2^{|I \cap K|} |P_I(S)| &= \sum_{I \subset \{1,2,\ldots,k+d\}} 2^{|I \cap K|} \\ &= \sum_{i=0}^{d} \sum_{j=0}^{k} \binom{k}{j} \binom{d}{i} 2^j \\ &= 3^k 2^d \end{aligned}$$

Now suppose that $|S| > 1$. We consider the same well-ordering $\prec$ on $\mathbb{Z}$ as considered in Section 2:

$$0 \prec 1 \prec -1 \prec 2 \prec -2 \prec 3 \cdots$$

and we denote by $<$ the regular ordering on either $\mathbb{Z}$ or $\mathbb{N}$. As in Section 2, for $a \in \mathbb{Z}$, we let a^+ denote the element of $\{a, a+1, a-1\}$ which is largest in the well-ordering $\prec$ and a^- the element of $\{a, a+1, a-1\}$ which is the smallest.

Let $z = (z_1, z_2, \ldots, z_{k+d}) \in S$ be a point which is a "corner point" of S. That is, for any $i \in \{1, 2, \ldots, k\}$, the point

$$(z_1, \ldots, z_{i-1}, z_i^+, z_{i+1}, \ldots, z_{k+d})$$

is not in S and for any $j \in \{k+1, k_2, \ldots, k+d\}$, the point

$$(z_1, \ldots, z_{j-1}, z_j + 1, z_{j+1}, \ldots, z_{k+d})$$

is not in S. We note that z exists because S is finite. Also note that, since S is compressed in every coordinate, for any $i \in \{1, 2, \ldots, k\}$ where $z_i \neq 0$,

$$(z_1, \ldots, z_{i-1}, z_i^-, z_{i+1}, \ldots, z_{k+d}) \in S$$

and

$$(z_1, \ldots, z_{i-1}, z_i^-, z_{i+1}, \ldots, z_{k+d}) \neq (z_1, \ldots, z_{i-1}, z_i, z_{i+1}, \ldots, z_{k+d})$$

Additionally, for any $j \in \{k+1, k+2, \ldots, d+k\}$ where $z_j \neq 0$,

$$(z_1, \ldots, z_{j-1}, z_j - 1, z_{j+1}, \ldots, z_{k+d}) \in S$$

and

$$(z_1, \ldots, z_{j-1}, z_j - 1, z_{j+1}, \ldots, z_{k+d}) \neq (z_1, \ldots, z_{j-1}, z_j, z_{j+1}, \ldots, z_{k+d})$$

By induction,

$$\partial \left(S \backslash \{z\}\right) = \sum_{I \subset \{1,2,\ldots,k+d\}} 2^{|I \cap K|} |P_I(S \backslash \{z\})|.$$

Define

$$Z_K = \{i \in \{1, 2, \ldots, k\} : z_i = 0\}$$

and

$$Z_D = \{j \in \{k+1, k+2, \ldots, k+d\} : z_j = 0\}$$

Then we note that the only points $(y_1, y_2, \ldots, y_{k+d})$ which are in ∂S but not in $\partial \left(S \backslash \{z\}\right)$ are of the form

$$y_i = \begin{cases} -1, 0, \text{ or } 1 & \text{if } i \in Z_K \\ y_i^+ & \text{if } i \in \{1, 2, \ldots, k\} \backslash Z_K \\ 0 \text{ or } 1 & \text{if } i \in Z_D \\ y_i + 1 & \text{if } i \in \{k+1, k+2, \ldots, k+d\} \backslash Z_D \end{cases}$$

Thus, we can see that there are a total of

$$3^{|Z_K|} 2^{|Z_D|}$$

points in ∂S which are not in $\partial \left(S \backslash \{z\}\right)$ so that

$$|\partial S| = |\partial S \backslash \{z\}| + 3^{|Z_K|} 2^{|Z_D|}$$

We also note that since S is compressed in every coordinate, the only $I \subset \{1, 2, \ldots, k+d\}$ for which $|P_I(S)|$ is larger than $|P_I(S \backslash \{z\})|$ must be of the form $I \subset Z_K \cup Z_D$. If $I \subset Z_K \cup Z_D$, then $|P_I(S)| = |P_I(S \backslash \{z\})| + 1$ and if $I \not\subset Z_K \cup Z_D$, then $|P_I(S)| = |P_I(S \backslash \{z\})|$. Thus, we see that

$$\begin{aligned}\sum_{I \subset \{1,2,\ldots,k+d\}} 2^{|I \cap K|} |P_I(S)| &= \sum_{I \subset \{1,2,\ldots,k+d\}} 2^{|I \cap K|} |P_I(S \backslash \{z\})| + \sum_{I \subset Z_K \cup Z_D} 2^{|I \backslash Z_D|} \\ &= \partial\left(S \backslash \{z\}\right) + \sum_{j=0}^{|Z_K|} \sum_{i=0}^{|Z_D|} \binom{|Z_D|}{i} \binom{|Z_K|}{j} 2^j \\ &= \partial\left(S \backslash \{z\}\right) + 3^{|Z_K|} 2^{|Z_D|}\end{aligned}$$

and hence our Theorem is proven.

□

We note that, in light of Theorem 1 and Theorem 2, we have the following Corollary:

Corollary 2. *Let X_n denote the set of all subsets $X \subset \mathbb{Z}^k$ of size n. Define the function $f : X_n \to \mathbb{R}$ as follows: for $X \in X_n$,*

$$f(X) = \sum_{I \subset \{1,2,\ldots,k\}} 2^{|I|} |P_I(X)|.$$

The minimum value of this function is achieved at an initial segment of $\mathbb{Z}^k$ of size n, according to the well-ordering $\prec$

Additionally, in light of Corollary 1 and Theorem 2, we have the following Corollary:

Corollary 3. *Let Y_n denote the set of all subsets $Y \subset \mathbb{N}^k$ of size n. Define the function $f : Y_n \to \mathbb{R}$ as follows: for $Y \in Y_n$,*

$$f(Y) = \sum_{I \subset \{1,2,\ldots,k\}} |P_I(Y)|.$$

The minimum value of this function is achieved at an initial segment of $\mathbb{N}^k$ of size n, according to the well-ordering $\prec_{\mathbb{N}}$

4 Final Remarks

The calculations in Section 3 allow us to understand some of the behavior of the boundaries of sets in our graph. These calculations rely on the technique which we call centralization. While compression is a more powerful technique, it requires that sets of minimal boundary be nested. The process of centralizing and computing boundaries does not require that sets of minimal boundary have any particular form. Perhaps these ideas can be used in other situations where the sets of minimal boundary are more complicated.

References

[1] Rudolf Ahlswede and Ning Cai. General edge-isoperimetric inequalities. I. Information-theoretical methods. *European J. Combin.*, 18(4):355–372, 1997.

[2] Rudolf Ahlswede and Ning Cai. General edge-isoperimetric inequalities. II. A local-global principle for lexicographical solutions. *European J. Combin.*, 18(5):479–489, 1997.

[3] Noga Alon, Itai Benjamini, and Alan Stacey. Percolation on finite graphs and isoperimetric inequalities. *Ann. Probab.*, 32(3A):1727–1745, 2004.

[4] A. Barvinok and A. Samorodnitsky. The distance approach to approximate combinatorial counting. *Geom. Funct. Anal.*, 11(5):871–899, 2001.

[5] B. Bollobás. Martingales, isoperimetric inequalities and random graphs. In *Combinatorics (Eger, 1987)*, volume 52 of *Colloq. Math. Soc. János Bolyai*, pages 113–139. North-Holland, Amsterdam, 1988.

[6] Béla Bollobás. Random graphs revisited. In *Probabilistic combinatorics and its applications (San Francisco, CA, 1991)*, volume 44 of *Proc. Sympos. Appl. Math.*, pages 81–97. Amer. Math. Soc., Providence, RI, 1991.

[7] Béla Bollobás and Imre Leader. An isoperimetric inequality on the discrete torus. *SIAM J. Discrete Math.*, 3(1):32–37, 1990.

[8] Béla Bollobás and Imre Leader. Compressions and isoperimetric inequalities. *J. Combin. Theory Ser. A*, 56(1):47–62, 1991.

[9] Béla Bollobás and Imre Leader. Edge-isoperimetric inequalities in the grid. *Combinatorica*, 11(4):299–314, 1991.

[10] Béla Bollobás and Imre Leader. Isoperimetric inequalities and fractional set systems. *J. Combin. Theory Ser. A*, 56(1):63–74, 1991.

[11] Thomas A. Carlson. The edge-isoperimetric problem for discrete tori. *Discrete Math.*, 254(1-3):33–49, 2002.

[12] L. H. Harper. Optimal numberings and isoperimetric problems on graphs. *J. Combinatorial Theory*, 1:385–393, 1966.

[13] L. H. Harper. *Global methods for combinatorial isoperimetric problems*, volume 90 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2004.

[14] Imre Leader. Discrete isoperimetric inequalities. In *Probabilistic combinatorics and its applications (San Francisco, CA, 1991)*, volume 44 of *Proc. Sympos. Appl. Math.*, pages 57–80. Amer. Math. Soc., Providence, RI, 1991.

[15] Michel Ledoux. *The concentration of measure phenomenon*, volume 89 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2001.

[16] John H. Lindsey, II. Assignment of numbers to vertices. *Amer. Math. Monthly*, 71:508–516, 1964.

[17] Jiří Matoušek. *Lectures on discrete geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002.

[18] Vitali D. Milman and Gideon Schechtman. *Asymptotic theory of finite-dimensional normed spaces*, volume 1200 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. With an appendix by M. Gromov.

[19] Gilles Pisier. *The volume of convex bodies and Banach space geometry*, volume 94 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1989.

[20] Oliver Riordan. An ordering on the even discrete torus. *SIAM J. Discrete Math.*, 11(1):110–127 (electronic), 1998.